

easy grow guide

geranium inspire

(F1 Pelargonium xhortorum)



Plug Production: 288 plugs

Sowing/Media:	Use a well-drained, disease-free, peat based plug medium with pH 6.0 - 6.2, EC <0.75mmhos. Cover seed with vermiculite
Germination Stage 1 & 2: (7 days)	Keep medium uniformly moist until seedlings are hooking above the covering, media temperature should be 72-75°F (22-24°C) but no higher as thermodormancy may occur, keep light levels <1500 f.c. Light is not needed for germination.
Germination Stage 3:	Media temperature can be dropped to 65-68°F (18-20°C), light levels should be <3000 f.c. HID lights can promote growth and faster flowering in periods of low light. Fertilize with 100-150ppm N from 15-5-15, 17-5-17 or 13-2-13, keep media pH at 6.0-6.5 no lower and EC <1.5mmhos.
Germination Stage 4:	Media temperatures can be lowered to 62-65°F (16-18°C), light levels should be maintained around 3000 f.c. Dry down between irrigations but avoid wilt. Fertilize with 100-150ppm N from 13-2-13 to tone seedlings. If needed, use sprays of Cycocel (250-750 ppm) or B-Nine (800 - 1500) + Cycocel (250 – 500 ppm) to control growth. It is best to run your own trials to avoid overdosing, as weather and cultural regimes can affect the requirements.

Growing On to Finish: Packs, 4" (10cm) pots

Media:	Use a well-drained, disease free, peat-based growing mix with pH 6.0 - 6.2 (no lower) and EC <1.5mmhos.
Temperatures:	Temperatures for rooting out after transplant should be 65-68°F (18-20°C) Temperatures for growing on can be lowered to 62-65°F (16-18°C)
Light:	Light levels should be 3000 - 5000 f.c. as a guide. HID lights can be used in low light periods to give more total light and encourage flowering.
Irrigation:	Practice a good wet/dry moisture cycle but avoid wilting. Geraniums do not like to be too wet but can't tolerate wilt either.
Fertilizer:	Feed 1-2 times per week with 150 – 200 ppm N from 15-5-15, 17-5-17, or 13-2-13, extra iron can be added if needed. Keep media pH 6.0 – 6.8, and media EC 1.25 – 1.75 mmhos.
Growth Regulators:	Use sprays of Cycocel (350-1500 ppm) or B-Nine (800 - 1500) + Cycocel (350 – 1000 ppm) to control growth. Bonzi sprays (2 – 5 ppm) can also be used once foliage covers the soil but do not drench with Bonzi as it has a strong effect on Geranium growth. It is best to run your own trials to avoid overdosing, as weather and cultural regimes can affect the requirements
Pests:	Aphids
Diseases:	Pythium, Botrytis, bacterial leafspot. Necrosis on lower leaves is likely to be caused by media pH <6.0, upper yellow leaves – high media pH >6.8 or low iron.

Plug Times:

288 plug:	4-5 weeks from sowing to transplant
------------------	-------------------------------------

Transplant to Finish:

Container	Plants per Container	Transplant to Finish	Total Crop Time
Packs	1 x plug	7-8 weeks	11-13 weeks
4" (10cm):	1 x plug	8-9 weeks	12-14 weeks

Crop times are based on optimum conditions. Alternative environmental conditions and cultural regimes can lengthen the crop times stated above.