

# Dune

## Marigold

Dune is a new, early blooming, intermediate sized African Marigold. Offering exceptional value for money, its fully double flowers have a unique flatter form, delivering excellent weather tolerance.

Seed Form	Clipped
Seed Count	12,750/oz - 450/g
Garden Height	10 - 12" (25 - 30cm)
Garden Spread	8 - 10" (20 - 25cm)
Flower Size	3 - 4" (8 - 10cm)



[www.floranova.com](http://www.floranova.com)



Dune Mixed MAR160



Gold MAR163



Orange MAR161



Yellow MAR162





# easy grow guide

## marigold dune

(OP *Tagetes erecta*)



### Plug Production: 512 or 288 plugs

<b>Sowing/Media:</b>	Use a well-drained, disease-free, peat based plug medium with pH 6.2-6.5 (avoid pH levels <6.0), EC <0.75mmhos. Cover seed with vermiculite
<b>Germination Stage 1: (3-5 days)</b>	Keep medium uniformly moist until seedlings are hooking above the covering, media temperature should be 68-72°F (20-22°C), keep light levels <1500 f.c.
<b>Germination Stage 2:</b>	Dry down slightly to avoid stretch, keep media temperature at 68-72°F (20-22°C), germination should be complete in 5-7 days.
<b>Germination Stage 3:</b>	Allow media to dry further between irrigations to improve rooting and control stretch, media temperature can be dropped to 65-68°F (18-20°C), light levels should be <3000 f.c. Fertilize with 100-150ppm N from 15-5-15, 17-5-17 or 13-2-13, keep media pH at 6.2-6.5 no lower and EC <1.5mmhos.
<b>Germination Stage 4:</b>	Practice good wet/dry moisture cycle, media temperatures can be lowered to 60-62°F (15-16°C), keep light levels <3000 f.c. Fertilize with 100-150ppm N from 13-2-13 to help tone the seedlings. If needed, use sprays of B-Nine (1500 – 2500 ppm), A-Rest (3 – 10 ppm), Bonzi (10 – 20 ppm), or Sumagic (2 – 7 ppm) to control growth. It is best to run your own trials to avoid overdosing, as weather and cultural regimes can affect the requirements.

### Growing On to Finish: Packs, 4.5 inch (11cm) pots

<b>Media:</b>	Use a well-drained, disease free, peat-based growing mix with pH 6.2-6.5 (no lower) and EC <1.5mmhos. pH levels below 6.0 can induce iron, manganese and sodium toxicity.
<b>Temperatures:</b>	Temperatures for rooting out after transplant should be 65-68°F (18-20°C) Temperatures for growing on can be lowered to 62-65°F (16-18°C)
<b>Light:</b>	Light levels should be 3000 - 5000 f.c. as a guide.
<b>Irrigation:</b>	Practice a good wet/dry moisture cycle.
<b>Fertilizer:</b>	Feed 1–2 times per week with 150 – 200 ppm N from 15-5-15, 17-5-17, or 13-2-13. Keep media pH 6.0 – 6.5, and media EC 1.25 – 1.75 mmhos (saturated paste). If plugs have buds and poor growth before transplanting, push growth with more NH <sub>4</sub> feed, such as 20-10-20, but watch out for lowering media pH too much.
<b>Growth Regulators:</b>	Use sprays of B-Nine (2500 – 5000 ppm), Bonzi (15 – 30 ppm), or Sumagic (5 – 10 ppm) as needed. Drench with Bonzi (2 – 5 ppm) or Sumagic (1 – 2 ppm) when plants are up to size and flowering. It is best to run your own trials to avoid overdosing, as weather and cultural regimes can affect the requirements
<b>Pests:</b>	Aphids, Whitefly, Thrips, Red Spider mite, Leafminer
<b>Diseases:</b>	Pythium, Botrytis, Alternaria leafspot. A preventative fungicide program is recommended. Leafspots (bacterial and Alternaria), bronzing or speckling on lower leaves is likely to be caused by media pH <6.0.

### Plug Times:

<b>512 Plug:</b>	3-4 weeks from sowing to transplant
<b>288 plug:</b>	4-5 weeks from sowing to transplant

### Transplant to Finish:

Container	Plants/Container	Transplant to Finish	Total Crop Time
<b>Packs</b>	1 x plug	6-7 weeks	9-11 weeks
<b>4.5 inch (11cm):</b>	1 x plug	7-8 weeks	10-12 weeks
<b>6 inch (15cm):</b>	1 x plug	9-10 weeks	12-14 weeks

Crop times are based on optimum conditions. Alternative environmental conditions and cultural regimes can lengthen the crop times stated above.