

Zenith

F1 Marigold



The unique Afro-French hybrid (triploid) from Floranova with more branching, more flowers, more heat tolerance and more rain tolerance. With no seed set, Zenith requires no dead heading to maintain its phenomenal flower power all Summer long.

Seed Form	Clipped, Coated
Seed Count	9,200/oz - 325/g
Garden Height	14" (35cm)
Garden Spread	14" (35cm)
Flower Size	2 - 3" (6 - 8cm)



Golden Yellow MAR651



Lemon Yellow MAR650




Orange MAR671



Red MAR695



Red & Gold MAR652



Yellow MAR681

Deep Orange MAR653

easy grow guide

triploid marigold zenith

(F1 Tagetes xhybridus)



Plug Production: 288 plugs

Sowing/Media:	Use a well-drained, disease-free, peat based plug medium with pH 6.0-6.2, EC <0.75mmhos. Cover seed with a thin layer of vermiculite, otherwise the seedlings will root into the vermiculite and make transplanting difficult.
Germination Stage 1: (3-5 days)	Keep medium uniformly moist until seedlings are hooking above the covering, media temperature should be 68-72°F (20-22°C), keep light levels <1500 f.c.
Germination Stage 2:	Dry down slightly to avoid stretch, keep media temperature at 68-72°F (20-22°C), germination should be complete in 5-7 days.
Germination Stage 3:	Allow media to dry further between irrigations to improve rooting and control stretch, but avoid wilting. Media temperature can be dropped to 65-68°F (18-20°C), light levels should be <3000 f.c. Fertilize with 100-150ppm N from 15-5-15, 17-5-17 or 13-2-13, keep media pH at 6.2-6.5 no lower and EC <1.5mmhos.
Germination Stage 4:	Practice good wet/dry moisture cycle, media temperatures can be lowered to 60-65°F (15-18°C), keep light levels <3000 f.c. Fertilize with 100-150ppm N from 13-2-13 to help tone the seedlings.
Growth Regulators:	Growth can be controlled naturally by good moisture management. If needed, use sprays of B-Nine (1500 – 2500 ppm), A-Rest (3 – 10 ppm), Bonzi (10 – 20 ppm), or Sumagic (2 – 7 ppm) to control growth. It is best to run your own trials to avoid overdosing, as weather and cultural regimes can affect the requirements.

Growing On to Finish: Packs, 4” (10.5cm) pots

Media:	Use a well-drained, disease free, peat-based growing mix with pH 6.2-6.5, EC 1.5mmhos
Temperatures:	Temperatures for rooting out after transplant should be 65-68°F (18-20°C). These can be lowered to 62-65°F (16-18°C) for growing on.
Light:	Light levels should be 3000 - 5000 f.c. as a guide.
Irrigation:	Practice a good wet/dry moisture cycle, avoiding wilt.
Fertilizer:	Feed 1–2 times per week with 150 – 200 ppm N from 15-5-15, 17-5-17, or 13-2-13. Keep media pH 6.2 – 6.5, and media EC 1.5 – 1.75 mmhos.
Growth Regulators:	Growth can be controlled naturally by good moisture management. Use sprays of B-Nine (2500 – 5000 ppm), Bonzi (15 – 30 ppm), or Sumagic (5 – 10 ppm) as needed. Drench with Bonzi (2 – 5 ppm) or Sumagic (1 – 2 ppm) when plants are up to size and flowering. It is best to run your own trials to avoid overdosing, as weather and cultural regimes can affect the requirements
Pests:	Aphids, Whitefly, Thrips, Spider mite, Leafminer
Diseases:	Pythium, Botrytis, Leafspots (bacterial and Alternaria), bronzing or speckling on lower leaves is likely to be caused by media pH <6.0.

Plug Times:

288 plug:	3-4 weeks from sowing to transplant
------------------	-------------------------------------

Transplant to Finish:

Container	Plants/Container	Transplant to Finish	Total Crop Time
Packs	1 x plug	4-5 weeks	7-9 weeks
4” inch (10.5cm):	1 x plug	5-6 weeks	8-10 weeks

Crop times are based on UK trials in optimum conditions. Alternative environmental conditions and cultural regimes can alter the crop times stated above.