



Sugar Rush Mixed WAL135



The Fall-flower Wallflower

Sugar Rush
F1 Wallflower

Sweetly scented and ultra-fast in production. Not requiring cool temperatures for flowering, this F1 hybrid Cheiranthus can be sold even in late Summer or early Autumn from as little as 9 weeks from sowing. Even during warm temperatures it still produces dark green healthy foliage and great natural branching. Once planted out it will withstand heavy frost and wintery conditions, making a nice bushy plant that will then re-flower in early Spring. Alternatively, it can be sold as a flowering plant in early Spring in as little as 14-16 weeks from sowing. The perfect addition for Autumn and Winter gardening. A versatile companion to Pansies!

(Sugar Rush is not available in all regions. Please refer to your pricelist for more information.)

<i>Seed Form</i>	Natural
<i>Seed Count</i>	18,400/oz - 650/g
<i>Garden Height</i>	8 - 10" (20 - 25cm)
<i>Garden Spread</i>	8 - 10" (20 - 25cm)
<i>Flower</i>	1 - 1½" (2 - 4cm)



Yellow WAL115



Primrose WAL112



Orange WAL127



Purple Bicolour WAL113



Red WAL124



Easy Grow Guide

Wallflower Sugar Rush

F1 *Cheiranthus cheirii*



Plug Production: 512 or 288 plugs

Sowing/Media:

Use a well-drained, disease-free, peat based plug medium with pH 5.5-5.8, EC <1.0 mmhos. Cover seed with coarse vermiculite

Germination Stages 1 & 2 (9-10 days):

Keep medium uniformly moist until hooking is observed, then dry off the vermiculite between irrigations. Media temperature should be 68-72°F (20-22°C), Light is not needed for germination but keep levels <1500 f.c.

Germination Stage 3:

Practice a good wet/dry cycle but avoid wilting, media temperature should be 62-65°F (17-18°C), light levels should be between 2500-3000 f.c. Fertilize every other watering with 100-150ppm N from 15-5-15, 17-5-17, or 13-2-13 keep media pH <6.5 and EC at 1.0-1.25 mmhos.

Germination Stage 4:

Dry down media between irrigations avoiding wilt and lower media temperature to 60-62°F (15-17°C), light levels should be around 3000 f.c. if possible. Fertilize as required but use more 13-2-13 if growing cooler than 60°F (15°C)

Growth Regulators:

A sprinch of Bonzi at 1ppm during emergence will prevent stretch and floppiness or alternatively a spray of Bonzi at 10ppm during stage 3 will have a similar effect. Weather conditions and cultural practices directly affect how much growth regulator to use, so it is recommended that you run your own trials.

Growing On to Finish: Packs, Jumbo Packs, 4" (10cm) pots

Media:

Use a well-drained, disease free, peat-based growing mix with pH 5.5-5.8, EC <1.5mmhos.

Temperatures:

Keep media temperature at 65°F(18°C) until roots have developed and then lower to 60°F(15°C) as an average daily temperature to grow on. Sugar Rush does not require vernalisation to induce flowering, it is the combination of cool temperatures (especially at night) and good light quality that gives the best results. They are happy to be grown alongside Pansies for fall and spring.

Light:

Keep light levels at 3000-5000 f.c but keep temperature cool. Sugar Rush has a facultative long day response so although it doesn't need long days to flower, it will flower faster when day length is 13 hours or more. Regardless of day length though, exposure to good light quality with ideal temperatures will reduce crop time when growing over winter, compared to when grown in cold, low light conditions (see below)

Irrigation:

Practice a good wet/dry moisture cycle avoiding extremes of wet or dry. Growing on the dry side is a good natural growth regulator and if the temperatures are cool it is important not to allow the plants to get too wet.

Fertilizer:

Fertilize 1-2 times per week with 150-200 ppm N from 15-5-15, 17-5-17, or 13-2-13, keep media pH <6.5 and media EC 1.5-2.0 mmhos.

Growth Regulators:

If conditions are favourable, the height of Sugar Rush can be successfully controlled by growing on the dry side. For high density crops grown in the fall at higher temperatures, you can control stretch with sprays of B-Nine (2000-3000 ppm) + Cycocel (1000 ppm) tank mixed, or sprays of Bonzi at 20ppm. It is always best to run your own trials as climatic conditions and culture can affect the results.

Pests:

Thrips, Two-Spotted Spider Mite, Red Spider Mite, Aphids

Diseases:

Botrytis, Sclerotinia, Powdery Mildew, Pythium, Rhizoctonia

Plug Times:

512/288 Plug: 4-5 weeks from sowing to transplant

Transplant to Finish: 1 plant per pot/cell

Sugar Rush is programmable and the finish times vary depending on the time of year they are sown, due to day length, temperature and light quality. The table below shows crop times from different sowing dates (n.b. these trials were run in the UK in natural conditions with no additional heat and light so the results may vary slightly when grown in a different climate).

Sowing Week	26	28	30	34	36	38	50	1
Temperature Range:	60-85°F 15-30°C	60-85°F 15-30°C	50-85°F 10-30°C	35-75°F 2-25°C	32-68°F 0-20°C	32-68°F 0-20°C	32-68°F 0-20°C	32-75°F 0-25°C
Weeks from transplant to finish:	5-6	5-6	5-6	8-9	14-15	14-15	15-16	14-15